



FOTIVDA[®]
(tivozanib) capsules

YOUR
FOTIVDA
HANDBOOK

The information you need to know
to keep moving forward on FOTIVDA

AVEO
ONCOLOGY
an LG Chem company

This handbook aims to empower patients by providing helpful information on various aspects of support and resources.

From treatment tips to ongoing education and support, we've got you covered.

“My oncologist said not to give up hope because there are always more tools in the toolkit.”

– Valeri, a Patient Ambassador for FOTIVDA® (tivozanib)

Valeri was compensated for her time and expense. Please note that Valeri's experience is her own and individual results may vary.

To hear Valeri's story and other patient stories you can visit www.FOTIVDA.com.

Please see the Important Safety Information on pages 14-15 or visit www.FOTIVDA.com



About FOTIVDA[®] (tivozanib)

FOTIVDA is a prescription medicine used to treat adults with advanced kidney cancer (advanced renal cell carcinoma or RCC) that has been treated with 2 or more prior medicines and has come back, or did not respond to treatment.

Here are some of the most important things you need to know about FOTIVDA:

- ✓ Proven effectiveness and established safety profile.
- ✓ Convenient oral dosing once daily, taken with or without food, for 21 days followed by a 7-day break.
- ✓ Personalized support program with a wide variety of services to help you throughout your treatment journey.

Get informed, be engaged, feel empowered

For more information about FOTIVDA, visit www.FOTIVDA.com

SELECT IMPORTANT SAFETY INFORMATION

Possible serious side effects can occur with FOTIVDA. Call or see your healthcare provider right away if you develop:

High blood pressure (hypertension). High blood pressure may be severe, including a sudden, severe increase in your blood pressure (hypertensive crisis) that can lead to death. Your healthcare provider should check your blood pressure after 2 weeks and at least monthly and may prescribe medicine to treat high blood pressure. You should check your blood pressure regularly and tell your healthcare provider if you have increased blood pressure or experience confusion, headaches, dizziness, chest pain, or shortness of breath.

Please see the Important Safety Information on [pages 14-15](#) or visit www.FOTIVDA.com



Not actual patient and physician

Dosing and Treatment Tips

Taking FOTIVDA[®] (tivozanib)

FOTIVDA is a prescription medicine used to treat adults with advanced kidney cancer (advanced renal cell carcinoma or RCC) that has been treated with 2 or more prior medicines and has come back, or did not respond to treatment. It is not known if FOTIVDA is safe and effective in children. Please keep out of reach of children.

SELECT IMPORTANT SAFETY INFORMATION

Possible serious side effects can occur with FOTIVDA. Call or see your healthcare provider right away if you develop:

High blood pressure (hypertension). High blood pressure may be severe, including a sudden, severe increase in your blood pressure (hypertensive crisis) that can lead to death. Your healthcare provider should check your blood pressure after 2 weeks and at least monthly and may prescribe medicine to treat high blood pressure. You should check your blood pressure regularly and tell your healthcare provider if you have increased blood pressure or experience confusion, headaches, dizziness, chest pain, or shortness of breath.

Please see the Important Safety Information on **pages 14-15** or visit **www.FOTIVDA.com**



Three weeks on treatment, one week off

- One capsule, once daily with water.
- With or without food.
- Every 21 days followed by a 7-day break.
- Take FOTIVDA® (tivozanib) as prescribed by your healthcare provider.
- Call your doctor right away if you take too much FOTIVDA or if you miss a dose.
- Do not stop taking FOTIVDA without speaking to your doctor.



Capsule shown not actual size.

Helpful reminders for taking FOTIVDA:

- ✓ Talk to your doctor about your treatment schedule and what to expect.
- ✓ Use a journal to track doses and potential symptoms.
- ✓ Take your medicine at the same time every day as part of your daily routine.
- ✓ Use reminders on your phone or calendar to take your daily dose.
- ✓ Stay organized and keep track of your appointments and refill reminders.
- ✓ Write down how you are feeling and any questions for your doctor.
- ✓ Build a support system to rely on for help such as family, friends, cancer advocates, and your healthcare team.

Please see the Important Safety Information on **pages 14-15** or visit www.FOTIVDA.com

Managing Potential Symptoms

Here are some helpful tips:

Fatigue

Fatigue is the feeling of being tired or weak. Symptoms of fatigue are normal during cancer treatment. Remember, everyone's experience is unique.

Here are some tips to help manage fatigue:



Conserve energy. Prioritize tasks, allocate energy wisely, and lean on family or friends for support.



Stay hydrated. Drink plenty of fluids.



Prioritize rest and sleep. Consider short naps during the day to recharge.



Physical activity. Engage in gentle exercise such as walking, stretching, or yoga.



Nutrition. Maintain a balanced diet. Focus on nutrient-rich foods to support your energy levels.

Diarrhea

Diarrhea is increased frequency of bowel movements, often associated with loose or watery stools.

Consult your doctor if the diarrhea persists or becomes severe. They may recommend an over-the-counter diarrhea medication or prescribe something more effective.

Tips to help with diarrhea:



Stay hydrated. Diarrhea can lead to dehydration. Drink plenty of clear liquids to keep yourself hydrated.



Avoid certain drinks. Caffeine and acidic drinks like orange juice can exacerbate diarrhea.



Space out your meals. Eat small, light meals throughout the day.

Please see the Important Safety Information on **pages 14-15** or visit **www.FOTIVDA.com**

FOTIVDA[®]
(tivozanib) capsules

Decreased appetite

Your treatment can interfere with your ability to drink enough fluids or eat enough food to maintain your weight and energy.

Here are some things you can do to help:



Stay active. Engaging in light physical activity before meals can help stimulate your appetite.



Avoid skipping meals. Even if you don't feel hungry, try to eat several small meals or snacks throughout the day instead of three large meals.



Socialize during meals. Sit down with family and friends and discuss topics other than your health.



Speak to a nutritionist. They may be able to recommend certain foods or liquids that will help you maintain your weight.

Nausea

Nausea is the feeling of being queasy or feeling that you may throw up.

Here are some ways to help manage nausea:



Stay hydrated. Drink small amounts of clear liquids such as water, ginger ale, or clear broth to help alleviate nausea.



Monitor your eating. Avoid spicy, fried, or highly sweet foods.



Eat small, frequent meals. Instead of large meals, opt for smaller, more frequent meals or snacks throughout the day. An empty stomach can worsen nausea.



Consult your doctor. If the nausea persists or worsens, they may recommend anti-nausea medications or other advice.

Remember that every patient is unique when it comes to symptoms, so it's essential that you seek personalized advice from your healthcare provider if you experience any of these symptoms.



Not actual physician and patient

Partnering with your Healthcare Team

Being diagnosed with cancer can be overwhelming, but you don't have to face it alone. Your healthcare team is there to support you every step of the way. Building a strong partnership with your healthcare providers ensures that you receive the best care possible. Here are some essential practices to enhance your collaboration:

Please see the Important Safety Information on **pages 14-15** or visit www.FOTIVDA.com



Before the Appointment:

Prepare Questions: Write down any questions or concerns you have about your cancer diagnosis, treatment options, symptoms, or overall care. Bring this list to your appointment.

Review Medical Records: Gather relevant medical records, including test results, previous treatments, and any changes in symptoms. Having this information readily available will help your healthcare team make informed decisions.

Understand Your Diagnosis: Research your specific cancer type, stage, and prognosis. Being informed empowers you to actively participate in discussions with your doctor.

Bring a Support Person: Consider bringing a family member or friend to appointments. They can take notes, ask questions, and provide emotional support.

Set Goals: Clarify your goals for treatment. Discuss what matters most to you—whether it's symptom relief, quality of life, or curative intent.

During the Appointment:

Active Listening: Pay attention to what your doctor says. Take notes or record the conversation (with permission) to review later.

Ask Questions: Don't hesitate to ask about treatment options, potential symptoms, and long-term effects. Seek clarification if something is unclear.

Discuss Preferences: Share your preferences regarding treatment. Consider factors like lifestyle, work, and family commitments.

Shared Decision-Making: Participate in shared decision-making. Understand the risks and benefits of each treatment option. Your voice matters.

Be Honest: Share any symptoms, concerns, or emotional challenges you're experiencing. Honesty helps your healthcare team tailor care to your needs.

Learn About Potential Symptoms: Discuss potential symptoms and how to manage them. Ask about strategies to minimize discomfort.

Follow-Up Plan: Understand the next steps—whether it's follow-up appointments, tests, or additional treatments. Clarify any uncertainties.



Remember that your healthcare team is there to support you. Effective communication ensures better outcomes and a more patient-centered experience.



Not actual physician and patient

Support and Resources

Financial support and resources

AVEO ACE is a comprehensive program dedicated to providing personalized support to patients and their loved ones throughout the FOTIVDA® (tivozanib) treatment journey.



CALL US AT:
1-833-FOTIVDA

(1-833-368-4832) M-F 8 AM to 8 PM ET



VISIT US AT:
WWW.FOTIVDA.COM



FAX US AT:
1-833-FOTIVDA

Please see the Important Safety Information on **pages 14-15** or visit **www.FOTIVDA.com**



How ACE can help:



- Understand your insurance coverage for FOTIVDA® (tivozanib).
- Learn more about your out-of-pocket costs for treatment.
- Identify financial support options that you may be eligible for.
- Access helpful resources and personalized nurse support.



Financial Assistance

- At AVEO ACE, we care about helping you access your FOTIVDA treatment. AVEO ACE Program Specialists can help explore financial support options that may help lower your out of pockets costs for FOTIVDA.
- If you are commercially insured...
 - You may qualify for the AVEO ACE Copay Assistance Program and pay as little as \$0 for each prescription.
 - Talk to your pharmacist or an AVEO ACE Specialist to see if you qualify, or visit www.fotivda.com/patient-support or an AVEO ACE Program.
 - Additional terms and conditions apply*
- If you are government insured...
 - An AVEO ACE Program Specialist can help you get connected with the appropriate third-party foundation for assistance.
- If you are uninsured, underinsured, or unable to afford your out-of-pocket costs, you may be eligible to receive FOTIVDA free of charge. An AVEO ACE Program Specialist can help see if you qualify for the AVEO ACE Patient Assistance Program.

Ongoing Education and Support



- Since it is important to take your FOTIVDA medication as prescribed, we have designed a unique nurse support program to help you stay on track.
- Patient support and education does not include medical advice. Your healthcare provider should always be your source for information about your condition and treatment.

*Coverage and reimbursement decisions are made by insurance companies following the receipt of claims. Not valid for prescriptions paid, in whole or in part, by Medicaid, Medicare, VA, DOD, TRICARE or other federal or state programs including any state pharmaceutical assistance programs. Not valid where taxed, restricted or prohibited by law. AVEO ACE reserves the right to modify or terminate the program at any time without notice. Savings provided by the program may vary depending on patients' out-of-pocket costs. Additional terms and conditions apply.

Please see the Important Safety Information on **pages 14-15** or visit www.FOTIVDA.com

More Support Resources For You

General cancer support

American Cancer Society

Cancer.org
800.227.2345

CancerCare

CancerCare.org

National Cancer Institute (NCI)

Cancer.gov
800.422.6237

Cancer Hope Network

Cancerhopenetwork.org
877-467-3638

Cancer support for advanced kidney cancer

KC CURE

Kccure.org

National Kidney Foundation

Kidney.org
855.653.2273

Kidney Cancer Association (KCA)

Kidneycancer.org
800.850.9132

Please see the Important Safety Information on **pages 14-15** or visit **www.FOTIVDA.com**

IMPORTANT SAFETY INFORMATION

FOTIVDA® (tivozanib) is a prescription medicine used to treat adults with advanced kidney cancer (advanced renal cell carcinoma or RCC) that has been treated with 2 or more prior medicines and has come back or did not respond to treatment.

It is not known if FOTIVDA is safe and effective in children.

Important Safety Information

Before taking FOTIVDA, tell your healthcare provider about all your medical conditions including, if you have high blood pressure, a history of heart failure, a history of blood clots in your veins or arteries (including stroke, heart attack, or change in vision), bleeding problems, thyroid problems, liver problems, an unhealed wound, if you plan to have surgery or have had recent surgery, or are allergic to FD&C No. 5 (tartrazine) or aspirin.

Tell your healthcare provider if you are pregnant or planning to be. FOTIVDA can harm your unborn baby. If you are able to become pregnant:

- Your healthcare provider should do a pregnancy test before you start treatment.
- Use effective birth control (contraception) during treatment and for 1 month after your last dose.
- Talk to your healthcare provider about birth control methods that may be right for you.
- Tell your healthcare provider right away if you become pregnant or think you might be pregnant.
- Do not breastfeed during treatment and for 1 month after your last dose of FOTIVDA.

Tell your healthcare provider about all the medicine you take and any new medicine. Taking FOTIVDA with certain other medicines may affect how FOTIVDA works.

FOTIVDA may cause serious side effects, including:

- **High blood pressure (hypertension).** High blood pressure may be severe, including a sudden, severe increase in your blood pressure (hypertensive crisis) that can lead to death. You should check your blood pressure regularly and tell your healthcare provider if you have increased blood pressure or experience confusion, headaches, dizziness, chest pain, or shortness of breath.
- **Heart failure.** Heart failure may be serious and sometimes lead to death. Your healthcare provider should check for symptoms of heart failure regularly, such as shortness of breath or swelling of your ankles.
- **Heart attack and blood clots in your veins or arteries.** Blood clots may be serious and sometimes lead to death. Tell your healthcare provider or get emergency medical help right away if you have, new chest pain or pressure, numbness or weakness on one side of your body, pain in your arms, back, neck or jaw, trouble talking, shortness of breath, sudden severe headache, vision changes, swelling in the arms or legs.

Please see the Important Safety Information on [pages 14-15](#) or visit www.FOTIVDA.com



- **Bleeding problems.** Bleeding may be serious and sometimes lead to death. Report or get medical help right away if you have, unusual bleeding from the gums, red or black stools (looks like tar), menstrual bleeding or vaginal bleeding that is heavier than normal, bruises that happen without a known cause or get larger, headaches, feeling dizzy or weak, bleeding that is severe or you cannot control, coughing up blood or blood clots, pink or brown urine, vomiting blood or your vomit looks like “coffee grounds,” unexpected pain, swelling, or joint pain
- **Protein in your urine.** Your healthcare provider should check your urine for protein before and during treatment.
- **Tear (perforation) in your stomach or intestines or an abnormal connection between two or more body parts (fistula).** Get medical help right away if you experience tenderness or pain in your stomach-area (abdomen) that is severe and does not go away.
- **Thyroid gland problems.** Your healthcare provider should do blood tests to check your thyroid gland function before and during your treatment and may prescribe medicine if you develop thyroid gland problems.
- **Risk of wound-healing problems.** Wounds may not heal properly during treatment. Tell your healthcare provider if you plan to have surgery before starting or during treatment, including dental surgery. You should stop taking FOTIVDA at least 24 days before planned surgery. Your healthcare provider should tell you when you may start taking FOTIVDA again after surgery.
- **Reversible Posterior Leukoencephalopathy Syndrome (RPLS).** RPLS is a condition that can happen. Tell your healthcare provider right away if you have headaches, seizures, confusion, blindness or changes in vision, or difficulty thinking.
- **Allergic reactions to tartrazine (FD&C Yellow No. 5).** FOTIVDA contains a dye called FD&C Yellow No. 5 (tartrazine) that may cause allergic-type reactions, including bronchial asthma, in certain people. This occurs most often in people who also are allergic to aspirin.

Common side effects include tiredness, diarrhea, decreased appetite, nausea, hoarseness, low levels of thyroid hormones, cough, mouth sores, decreased blood levels of salt (sodium) and phosphate, increased levels of lipase in the blood.

Other side effects include vomiting and weakness or lack of energy.

FOTIVDA may cause fertility problems in males and females, which may affect your ability to have a child.

Your healthcare provider may change your dose, temporarily stop, or permanently stop treatment with FOTIVDA if you have certain side effects.

These are not all the possible side effects of FOTIVDA.

To report SUSPECTED ADVERSE REACTIONS, contact AVEO Pharmaceuticals, Inc. at 1-833-FOTIVDA (1-833-368-4832) or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see the Patient Information in the full Prescribing Information for FOTIVDA® (tivozanib).

Please see the Important Safety Information on **pages 14-15** or visit **www.FOTIVDA.com**

